



Baker Research Online
<http://library.bakeridi.edu.au>

This is the postprint version of the work. It is the manuscript that was accepted by the journal following peer review. It does not include the publisher's layout and pagination.

Shi Z, Zhen S, Zimmet PZ, Zhou Y, Zhou Y, Magliano DJ, Taylor AW. Association of impaired fasting glucose, diabetes and dietary patterns with mortality: a 10-year follow-up cohort in Eastern China. *Acta Diabetol* 2016;53(5):799-806.

<http://hdl.handle.net/11187/2686>

Copyright © Springer Verlag. This file is for personal use. Further distribution is not permitted.
The final publication is available at link.springer.com

Association of impaired fasting glucose, diabetes and dietary patterns with mortality: a 10-year follow-up cohort in Eastern China

Zumin Shi ^{1,2*}, Shiqi Zhen ¹, Paul Z Zimmet ^{3,4}, Yonglin Zhou ¹, Yijing Zhou ¹,
Dianna J Magliano ³, Anne W Taylor ²

¹ Department of Nutrition and Foodborne Disease Prevention, Jiangsu Provincial Centre for Disease Control and Prevention, China

² Discipline of Medicine, University of Adelaide, Level 7, SAHMRI, North TCE, Adelaide, Australia, 5000

³ BakerIDI Heart and Diabetes Institute, Melbourne, Australia

⁴ Department of Medicine, Monash University, Melbourne, Australia

Short title: Diabetes and mortality

Word count: 2520

Number of Tables: 2

Number of Figures: 3

Corresponding Author: A/Professor Zumin Shi

Postal address; Discipline of Medicine, University of Adelaide, Level 7, SAHMRI, North Terrace, Adelaide, Australia, 5005

Phone: +61 8 8313 1188

Fax: +61 8 8313 1228

Email: Zumin.shi@adelaide.edu.au

Abstract

Aims: To examine the association between impaired fasting glucose (IFG)/type 2 diabetes and mortality as well as to explore any interactions with dietary intake patterns in a Chinese population.

Methods: We followed 2849 Chinese adults aged 20 years and older for 10 years. Fasting plasma glucose was measured at baseline in 2002. Dietary patterns were constructed using factor analysis. Hazard ratios (HRs) and 95%CI were calculated by Cox proportional hazards analysis (all-cause mortality) and competing risks regression (cardiovascular disease (CVD)).

Results: Of the 2849 participants, 102 had diabetes and 178 had impaired fasting glucose (IFG) at baseline. We documented 184 deaths (70 CVD deaths) during 27,914 person-years of follow-up. Diabetes was associated with death from all-causes (HR 2.69, 95% confidence interval (CI) 1.62 - 4.49) after adjusting for socio-demographic and lifestyle factors. Diabetes had a HR of 1.97 (95% CI 0.84-4.60) for CVD death. IFG had 83% increased risk of all-cause mortality. Among those with low and high intake of a vegetable rich dietary pattern, the HR of IFG/diabetes for all-cause mortality were 3.25 (95%CI 1.95-5.44) and 1.38 (95% CI 0.75-2.55) (p for interaction 0.019), respectively.

Conclusions: Diabetes and IFG are associated with a substantial increased risk of death in Chinese adults. Dietary patterns associated with a high intake of vegetable was associated with a decrease in the risk of mortality for those with IFG/diabetes.

Key words: Diabetes, mortality, Chinese, cohort study, dietary pattern

Introduction

The world is facing a diabetes epidemic with 415 million people living with diabetes [1]. China is one of the countries with the highest prevalence of diabetes [2, 3]. It is well known that people with diabetes have a lower life expectancy. In Western countries, numerous data show that diabetes increases the risk of mortality, especially from cardiovascular disease (CVD) [4-8]. However, studies in Asia in this area are limited [9]. Several studies have been conducted among the Chinese population outside the mainland of China [10-13]. In China, most of the studies on the association between diabetes and mortality are hospital-based clinical studies or from chronic disease registries [14, 15]. There is only one general population based study- the Da Qing IGT and Diabetes Study, which was conducted in northern China, [16] and was based on a trial population.

White rice has a high glycemic index and a high intake of refined rice increases the risk of diabetes [17]. Compared with wheat as the staple food, rice intake is inversely associated with weight gain and high blood pressure [18]. Findings from Japan suggest that high rice intake is not associated with the risk of all-cause mortality in the general population [19]. However, the interaction between rice intake, rice based dietary patterns and IFG/diabetes in relation to mortality has not been studied in Japan or elsewhere. The aim of this study was to assess the interaction between IFG/diabetes and 10-year mortality as well as to explore any interactions with dietary intake patterns among Chinese adults in southeast China.

Methods

Study population

The Jiangsu Nutrition Study (JIN) is an ongoing cohort study investigating the association of nutrition and other factors with the risk of non-communicable chronic disease [20, 21]. The sample was based on a subsample of the Chinese national nutrition and health survey representing Jiangsu province and the year 2002 was used as a baseline. The rural sample was selected from six counties (Jiangyin, Taichang, Shuining, Jurong, Sihong and Haimen). The urban sample was selected from the capital cities of the two prefectures (Nanjing and Xuzhou). In 2002, 2849 adults aged 20 and above had fasting blood samples analysed for plasma glucose and haemoglobin at the study sites.

The study was conducted according to the guidelines laid down in the Declaration of Helsinki, and all procedures involving human subjects/patients were approved by Jiangsu Provincial Centre for Disease Control and Prevention. Written informed consent was obtained from all the participants.

Data collection and measurements

Participants were interviewed at their homes by health workers using a standard questionnaire.

Categories of glucose metabolism We defined diabetes as fasting plasma glucose (FPG) >7.0 mmol/l or having known diabetes (self-reported doctor diagnosed).

Impaired fasting glucose (IFG) was defined as FPG between 5.6 and 6.9 mmol/l without known diabetes. FPG <5.6 mmol/l was defined as normal fasting glucose (NFG) [22].

Dietary measurements –Frequency of use and quantity of intake of 33 food groups and beverages during the past year was assessed using a validated food frequency

questionnaire (FFQ) [23]. Dietary patterns were constructed using factor analysis. Four dietary patterns were identified. Factor 1 ('macho') was characterised by various kinds of animal foods and alcohol. Factor 2 ('traditional') loaded heavily on rice and fresh vegetables and inversely on wheat flour. Factor 3 ('sweet tooth') had high loadings of cake, milk, yoghurt and beverages. Factor 4 ('vegetable rich' pattern) had high intake of whole grains, fruits, vegetables, wheat flour, milk, eggs and fish but a low intake of rice (Supplement Figure 1). Participants were assigned pattern-specific factor scores. Scores for each pattern were calculated as the sum of the products of the factor loading coefficients and standardized weekly intake of each food associated with that pattern. Detailed description of the dietary pattern identification has been published elsewhere [23]. In the analyses, each dietary pattern was dichotomised as low and high intake based on factor scores.

In addition to the FFQ based food intake calculation, food intake including rice intake were assessed using a 3-day weighed food record (WFR) which recorded all foods consumed by each individual, on three consecutive days (including one weekend) (N=2832). At the beginning and end of the 3-day survey, health workers weighed all the food stocked in the household. Each day, all purchases, home production, and processed snack foods were weighed and recorded. Food intakes of each individual in the household were recorded in detail each day. During the interview, the health workers would check any intake value for a particular food that fell below or above the usual intake value by the population in the region. For the current study, rice intake (presented as raw rice, g/d) for each individual was calculated based on the 3-day records. We did not consider under- and over-reporting of energy intake as an issue of concern because any unreliable data was checked by the health workers during the survey. Food consumption data were analysed using the Chinese Food

Composition Table [24] . In the analysis, rice intake was divided by total energy intake and further recoded into tertiles.

Covariates Cigarette smoking was assessed by asking frequency of daily cigarette smoking. Alcohol consumption was assessed by asking the frequency and amount of alcohol/wine intake. Information on physical activity was collected using a validated physical activity questionnaire covering a time period of one year [25]. Active commuting (walking or cycling to and from work) was categorized into three groups: none, 1- 30 min per day and more than 30 min per day. Daily leisure time physical activity was classified into two categories (yes, or no). Sedentary activity (reading, watching TV, using computer, and playing video games) was categorised into three groups (<1 hour/d, 1-2 hours/d and >2 hours/d). Occupation was recoded into manual or non-manual based on a question with 12 occupational categories. Among people with known diabetes/dyslipidaemia, whether dietary change was used to manage the condition was asked. Hypertension was defined as systolic blood pressure above 140 mmHg and/or diastolic blood pressure above 90 mmHg, or using antihypertensive drugs. Overweight was defined as BMI ≥ 24 kg/m².

Death ascertainment The underlying cause of mortality was defined according to the World Health Organization International Classification of Disease, 10th revision (ICD-10). Information of death was collected in 2012 by household visit as well as by linking with the death registry database in the local Centre for Disease Control and Prevention. Thus, the identification of death was virtually complete. CVD mortality included ICD-10 codes I00–99. Cancer mortality was defined as ICD-10 C00–97.

Statistical analysis

The chi square test was used to compare differences in categorical and ANOVA in continuous variables. For each participant, person-years of follow-up were calculated from the date of baseline survey to the date of death or the date of last follow-up (1 December 2012), whichever came first. The association between diabetes/IFG and the risk of CVD was analysed using competing risks regression (*stcrreg* syntax in Stata) and the association between diabetes/IFG and all-cause mortality was analysed using Cox proportional hazard models, adjusting for multiple covariates. Two models assessed the association between diabetes/IFG and mortality. The first model controlled for age (continuous) and gender; the second model further adjusted for sociodemographic and lifestyle factors, chronic diseases (anaemia, BMI, and hypertension), and intake of energy, fat and fibre. As the sample size in the full model was 2757 (96.8% of the whole sample), we did not impute the missing data. Due to the low number of deaths, we also modelled a combined diabetes/IFG to reflect abnormal glucose tolerance. The proportional hazards assumption in the Cox model was assessed with graphical methods and with models including time-by-covariate interactions. In general, all proportionality assumptions were appropriate.

Interactions between diabetes/IFG and lifestyle factors (dietary patterns, rice intake, smoking, alcohol drinking, and physical activity), sociodemographic factors (sex, occupation, residence, and income), and overweight were conducted by adding a multiplicative term in the fully adjusted models. Results were presented graphically.

As all the members in the household were invited to participate in the study, we used a cluster-robust variance estimator [stata command: *vce(cluster clustvar)*] to account for the clustering at the household level in the estimation of the variance.

This estimator relaxes the assumption of independence of the observations and produces the correct standard errors. Statistical significance was considered when $P <$

0.05 (two-sided). All analyses were performed using Stata 14 (Stata Corp., College Station, TX, USA).

Results

Of the 2849 participants, 102 and 178 had diabetes and IFG respectively at baseline. Their mean age was 47.0(SD 14.5) years. **Table 1** shows the sample characteristics by glucose metabolism categories. About half of those with diabetes reported dietary change to manage chronic disease. However, this figure was only 6% among those with IFG. The mean intake of raw rice was above 100 g per 1000 kcal across all categories of glucose metabolism.

We documented 184 deaths (70 CVD deaths and 63 cancer deaths) during 27,914 person-years of follow-up. The overall all-cause mortality was 6.6 per 1000 person-years. The crude all-cause mortality was 5.8, 12.4 and 17.6 per 100 person-years among those with NFG, IFG and diabetes, respectively. **Figure 1** shows the age adjusted cumulative mortality by baseline glucose metabolism categories.

Diabetes (vs. NFG) was associated with death from all-causes (HR 2.69, 95% confidence interval (CI) 1.62-4.49) after adjusting for sociodemographic and lifestyle factors (**Table 2**). Additional adjustment for diet change slightly attenuated the HR for all-cause mortality to 2.59 (95%CI 1.46-4.59) (data not shown). Participants with diabetes at baseline had a HR of 1.97 (95% CI 0.84-4.60) for CVD death. Compared with NFG, IFG had increased risk of all-cause mortality (HR 1.83, 95%CI 1.13-2.97). When IFG/diabetes was combined, the HRs associated with all cause and CVD mortality were 2.12 (1.44-3.12) and 1.68 (0.87-3.24), respectively (**Table 2**)

Interactions

A vegetable rich dietary pattern was inversely associated with all-cause mortality (for 1 SD increase of factor score, HR for all-cause mortality was 0.82 (95%CI 0.67-0.99)) in the fully adjusted model. There was a significant interaction between vegetable rich

dietary pattern and IFG/diabetes in relation to all-cause mortality (p for interaction 0.019) (**Figure 2**). The HRs of IFG/diabetes for all cause-mortality were 3.25 (95%CI 1.95-5.44) and 1.38 (95% CI 0.75-2.55) among those with a low and high intake of vegetable rich dietary pattern, respectively. No interaction between other dietary patterns and IFG/diabetes was found. There was a trend of increasing HRs of IFG/diabetes for all-cause mortality with the increase of rice intake, although p for interaction was not statistically significant (using rice intake as a continuous variable, p for interaction 0.199). With the increase of income, the HRs of IFG/diabetes for all-cause mortality decreased (p for interaction 0.193). Among those with low income, IFG/diabetes had a HR of 2.52 (95%CI 1.43-4.43) for all-cause mortality.

There was no significant interaction between diabetes and residence or overweight in relation to mortality.

Discussion

In this general population based prospective study, we found that both diabetes and IFG are associated with a substantial increased risk of all-cause mortality in Chinese adults. Compared with those with NFG, all-cause mortality risk in those with diabetes increased two and half times. Furthermore, there was a ~80% increased mortality from all-cause among those with IFG. We also detected an interaction between diabetes and vegetable rich dietary pattern in relation to mortality. In those with diabetes who had a low intake of vegetable rich pattern had an increased risk of mortality compared to those who had a high intake of the pattern. However, overweight and residence (urban/rural) were not associated with any increase in risk of mortality for those with diabetes.

A two to three times excess risk of mortality has often been found among those with diabetes as compared with those without diabetes [10, 16]. Reductions in all-cause mortality among those with diabetes have occurred over time [26]. The latest data from Sweden suggest that the overall excess risk of death had dropped to a historically low level of 15% [8]. In the Da Qing Study, the age-adjusted relative risk of death from all causes in those with diabetes was three times higher compared with those without diabetes [16]. However, the Da Qing study started in 1980's and over the past three decades there has been substantial development in the primary and secondary prevention of diabetes. Surprisingly, the excess risk of mortality in our study remained as high as 2.69 times. It may underline the poor control of glycemic level [3] as well as other risk factors among persons with diabetes in China. In our study, the prevalence of smoking was similar to the national average [27], A very low proportion of patients reported undertaking leisure time physical activity.

Furthermore, the consumption of refined wheat rice was high. In the IFG and diabetes

groups, the raw rice consumption was 117 g and 106 g per 1000 kcal, respectively. These unhealthy lifestyles may lead to the poor control of glycemic level and contribute to a high proportion of mortality among people with IFG/diabetes in China. The interaction between diabetes and income suggest that those in the disadvantaged population should be given priority for intervention.

The interaction between diabetes and vegetable rich dietary pattern is consistent with current knowledge about the beneficial effects of fruit, vegetable, and whole grain. The pattern is inversely associated with rice intake. It is also consistent with current knowledge about the relationship between rice intake and diabetes [17]. Rice has a high glycemic index and has been previously found to be associated with an increased risk of hyperglycemia in the cohort [18]. In general, a diet with a high glycemic index impairs glycemic control. It has been shown that poor glycemic control increases the risk of mortality [8].

Overall, our study suggests a beneficial effect of engaging in physical activity and a high intake of vegetable rich dietary pattern among those with diabetes. Because the prevalence of favourable lifestyle factors in the population is low, there appears an opportunity that lifestyle modification will be able to reduce a substantial number of premature deaths among those with diabetes.

Several limitations in our study should be acknowledged. Firstly, although we had 5-year follow-up, due to the high attrition rate, we decided not to incorporate the data at follow up. During a 5-year follow-up, 125 developed IFG/diabetes among 1056 participants with NFG at baseline. Therefore, our study may underestimate the risk of mortality among those with diabetes or IFG. Secondly, due to the small number of deaths, we could not further analyse mortality due to specific CVD outcomes due to

lack of statistical power. We also note that the number of deaths among those with diabetes was low and this has led to some uncertainty in our findings, especially for CVD mortality. However, despite this, the magnitude of the HR of diabetes for all cause and CVD mortality are in line with several well-established studies suggesting that our results are not likely to be due to chance. Education was positively with the intake of vegetable rich dietary pattern in the sample, but income was inversely associated with vegetable rich dietary pattern (data not shown). Thus the interaction between vegetable rich dietary pattern and IFG/diabetes could be confounded by socioeconomic status. Although we have adjusted for education and income in the multivariable model, residual confounding is still possible. Furthermore, dietary intake was not continuously measured during follow-up. The eating habits could change during the ten-year period. Finally, in our study, diabetes was based on fasting glucose and self-reported known diabetes. It is known that the prevalence of undiagnosed diabetes is high in China. An underestimation of the prevalence of diabetes is highly likely and this may affect the outcomes of the study.

The strength of the study is the face to face based interview by well-trained health workers at baseline as well as the use of 3-day WFR to measure energy and nutrients intake. The study included both urban and rural participants. It can be generalized in the study population to the province with over 70 million residents. However, whether the findings can be generalized in the whole Chinese population is not clear. In conclusion, diabetes and IFG are associated with an increased risk of death in Chinese adults. A high intake of vegetable rich dietary pattern is associated with decreased risk of mortality for those with diabetes. There is no interaction between the rice based traditional dietary pattern and IFG/diabetes.

Acknowledgments

The research was supported by The University of Adelaide and Jiangsu Provincial Centre for Disease Control and Prevention.

Author's contributions

The study was conceived and the statistical analysis carried out by Zumin Shi. Zumin Shi had full access to the data and drafted the manuscript. Critical revision of the manuscript for important intellectual content: Shiqi Zhen, Paul Z Zimmet, Yonglin Zhou, Yijing Zhou, Dianna J Magliano, Anne W Taylor.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflicts of interest.

Ethical standard All procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1975, as revised in 2008 (5).

Human and animal rights This study was approved by the Jiangsu Provincial Center for Disease Control and Prevention Institutional Review Board.

Informed consent Informed consent was obtained from all participants for being included in the study.

Reference

1. International Diabetes Federation. IDF Diabetes Atlas. 7th edition 2015.
2. Zimmet PZ, Magliano DJ, Herman WH, Shaw JE. Diabetes: a 21st century challenge. *Lancet Diabetes Endocrinol.* 2014;2(1):56-64. doi:10.1016/S2213-8587(13)70112-8.
3. Xu Y, Wang L, He J, Bi Y, Li M, Wang T et al. Prevalence and control of diabetes in Chinese adults. *JAMA.* 2013;310(9):948-59. doi:10.1001/jama.2013.168118.
4. Barr EL, Zimmet PZ, Welborn TA, Jolley D, Magliano DJ, Dunstan DW et al. Risk of cardiovascular and all-cause mortality in individuals with diabetes mellitus, impaired fasting glucose, and impaired glucose tolerance: the Australian Diabetes, Obesity, and Lifestyle Study (AusDiab). *Circulation.* 2007;116(2):151-7. doi:10.1161/CIRCULATIONAHA.106.685628.
5. Emerging Risk Factors C, Seshasai SR, Kaptoge S, Thompson A, Di Angelantonio E, Gao P et al. Diabetes mellitus, fasting glucose, and risk of cause-specific death. *N Engl J Med.* 2011;364(9):829-41. doi:10.1056/NEJMoa1008862.
6. Huxley R, Barzi F, Woodward M. Excess risk of fatal coronary heart disease associated with diabetes in men and women: meta-analysis of 37 prospective cohort studies. *BMJ.* 2006;332(7533):73-8. doi:10.1136/bmj.38678.389583.7C.
7. Hansen MB, Jensen ML, Carstensen B. Causes of death among diabetic patients in Denmark. *Diabetologia.* 2012;55(2):294-302. doi:10.1007/s00125-011-2383-2.
8. Tancredi M, Rosengren A, Svensson AM, Kosiborod M, Pivodic A, Gudbjornsdottir S et al. Excess Mortality among Persons with Type 2 Diabetes. *N Engl J Med.* 2015;373(18):1720-32. doi:10.1056/NEJMoa1504347.
9. Woodward M, Zhang X, Barzi F, Pan W, Ueshima H, Rodgers A et al. The effects of diabetes on the risks of major cardiovascular diseases and death in the Asia-Pacific region. *Diabetes Care.* 2003;26(2):360-6.

10. Magliano DJ, Soderberg S, Zimmet PZ, Cartensen B, Balkau B, Pauvaday V et al. Mortality, all-cause and cardiovascular disease, over 15 years in multiethnic mauritius: impact of diabetes and intermediate forms of glucose tolerance. *Diabetes Care*. 2010;33(9):1983-9. doi:10.2337/dc10-0312.
11. Ma S, Cutter J, Tan CE, Chew SK, Tai ES. Associations of diabetes mellitus and ethnicity with mortality in a multiethnic Asian population: data from the 1992 Singapore National Health Survey. *Am J Epidemiol*. 2003;158(6):543-52.
12. Tseng CH. Mortality and causes of death in a national sample of diabetic patients in Taiwan. *Diabetes Care*. 2004;27(7):1605-9.
13. Lin WH, Hsu CH, Chen HF, Liu CC, Li CY. Mortality of patients with type 2 diabetes in Taiwan: a 10-year nationwide follow-up study. *Diabetes Res Clin Pract*. 2015;107(1):178-86. doi:10.1016/j.diabres.2014.09.021.
14. Zhu M, Li J, Li Z, Luo W, Dai D, Weaver SR et al. Mortality rates and the causes of death related to diabetes mellitus in Shanghai Songjiang District: an 11-year retrospective analysis of death certificates. *BMC Endocr Disord*. 2015;15:45. doi:10.1186/s12902-015-0042-1.
15. Wei ZH, Xie XQ, Wei JM. Impact of ageing and related factors on death rate of diabetes mellitus in Beijing. *Zhonghua Yu Fang Yi Xue Za Zhi*. 2005;39(4):277-9.
16. An Y, Zhang P, Wang J, Gong Q, Gregg EW, Yang W et al. Cardiovascular and All-Cause Mortality Over a 23-Year Period Among Chinese With Newly Diagnosed Diabetes in the Da Qing IGT and Diabetes Study. *Diabetes Care*. 2015;38(7):1365-71. doi:10.2337/dc14-2498.
17. Hu EA, Pan A, Malik V, Sun Q. White rice consumption and risk of type 2 diabetes: meta-analysis and systematic review. *BMJ*. 2012;344:e1454. doi:10.1136/bmj.e1454.

18. Shi Z, Taylor AW, Hu G, Gill T, Wittert GA. Rice intake, weight change and risk of the metabolic syndrome development among Chinese adults: the Jiangsu Nutrition Study (JIN). *Asia Pac J Clin Nutr*. 2012;21(1):35-43.
19. Eshak ES, Iso H, Yamagishi K, Kokubo Y, Saito I, Yatsuya H et al. Rice consumption is not associated with risk of cardiovascular disease morbidity or mortality in Japanese men and women: a large population-based, prospective cohort study. *Am J Clin Nutr*. 2014;100(1):199-207. doi:10.3945/ajcn.113.079038.
20. Shi Z, Zhou M, Yuan B, Qi L, Dai Y, Luo Y et al. Iron intake and body iron stores, anaemia and risk of hyperglycaemia among Chinese adults: the prospective Jiangsu Nutrition Study (JIN). *Public Health Nutr*. 2010;13(9):1319-27. doi:S1368980009991868 [pii] 10.1017/S1368980009991868.
21. Shi Z, Hu X, Yuan B, Pan X, Meyer HE, Holmboe-Ottesen G. Association between serum ferritin, hemoglobin, iron intake, and diabetes in adults in Jiangsu, China. *Diabetes Care*. 2006;29(8):1878-83. doi:29/8/1878 [pii] 10.2337/dc06-0327.
22. American Diabetes A. Diagnosis and classification of diabetes mellitus. *Diabetes Care*. 2010;33 Suppl 1:S62-9. doi:10.2337/dc10-S062.
23. Shi Z, Hu X, Yuan B, Hu G, Pan X, Dai Y et al. Vegetable rich food pattern is related to obesity in China. *Int J Obes (Lond)*. 2008 32:975-84.
24. Yang Y. *Chinese Food Composition Table 2004*. Beijing: Peking University Medical Press; 2005.
25. Ma G, Luan D, Liu A, Li Y, Cui Z, Hu X et al. The analysis and evaluation of a physical activity questionnaire of Chinese employed population. *Ying Yang Xue Bao*. 2007;29:217-21.

26. Preis SR, Hwang SJ, Coady S, Pencina MJ, D'Agostino RB, Sr., Savage PJ et al. Trends in all-cause and cardiovascular disease mortality among women and men with and without diabetes mellitus in the Framingham Heart Study, 1950 to 2005. *Circulation*. 2009;119(13):1728-35. doi:10.1161/CIRCULATIONAHA.108.829176.
27. Li Q, Hsia J, Yang G. Prevalence of Smoking in China in 2010. *N Engl J Med*. 2011;364(25):2469-70. doi:doi:10.1056/NEJMc1102459.

Figure legends

Figure 1 Age adjusted cumulative incidence of all-cause mortality according to glucose categories: the Jiangsu Nutrition Study.

Figure 2 Adjusted HR of IFG/diabetes (*vs.* NFG) for all-cause mortality according to sociodemographic and lifestyle factors.

Except for stratification variables, all models adjusted for age, gender, , intake of energy, fat and fibre, smoking (0, 1-19, ≥ 20 cigarettes/day), alcohol drinking (no, 1-2 times/wk, 3-4 times/wk, daily), active commuting (no, 1-29 minutes/day, ≥ 30 minutes/day), leisure time physical activity (no, 1-29 minutes/day, > 30 minutes/day), sedentary activity (<1 hrs/day, 1-1.9 hrs/day, 2-2.9 hrs/day, ≥ 3 hrs/day), education (low, medium, high), occupation (manual/non-manual), BMI (linear and square terms) and hypertension.

P values were for interactions between IFG/diabetes and sociodemographic and lifestyle factors. Using rice intake as continuous variable, p for rice and IFG/diabetes interaction was 0.199.

Supplement Figure 1 Factor loadings of dietary patterns

Factor loadings $<|0.3|$ were not shown.

Table 1 Baseline characteristics of sample according to categories of glucose metabolism

	NFG	IFG	Diabetes	p-value
N	2569	178	102	
Age (years)	46.3 (14.4)	52.5 (13.8)	55.8 (12.3)	<0.001
BMI (kg/m ²)	23.3 (3.4)	25.0 (3.6)	25.8 (3.8)	<0.001
Overweight	37.2	58.4	65.7	<0.001
Men	45.8	47.2	46.1	0.94
Low income	33.4	25.7	18.6	<0.001
Low education	47.8	48.0	44.1	0.95
Manual job	55.1	40.4	23.5	<0.001
Hypertension	26.6	49.4	54.9	<0.001
Urban dwellers	23.6	30.9	50.0	<0.001
Smoking	27.2	25.0	29.7	0.69
Alcohol drinking	24.7	24.2	22.8	0.89
Having exercise	8.9	12.9	31.4	<0.001
Diet change to manage				
NCD	3.6	6.2	49.0	<0.001
Intake of food and nutrients				
Rice (g/1000 kcal)	109.6 (60.2)	117.0 (61.2)	106.3 (53.2)	0.23
Energy (kcal/d)	2371.2 (684.7)	2193.5 (631.1)	2108.6 (766.4)	<0.001
Fat (g/d)	81.0 (37.1)	78.6 (36.4)	79.9 (38.4)	0.695
Protein (g/d)	72.8 (23.6)	70.2 (22.2)	71.8 (29.3)	0.371
Carbohydrate (g/d)	328.2 (107.2)	295.3 (91.4)	267.9 (108.0)	<0.001
Dietary pattern scores				
Macho	-0.00 (0.99)	-0.01 (1.24)	0.02 (0.91)	0.967
Traditional	-0.02 (1.01)	0.09 (0.89)	0.37 (0.98)	<0.001
Sweet tooth	-0.01 (1.00)	0.12 (1.06)	0.08 (0.94)	0.179
Vegetable rich	-0.02 (0.96)	-0.02 (1.23)	0.56 (1.38)	<0.001

Values are % or mean (SD)

Table 2 HRs (95 CIs) for all-cause and CVD mortality according to categories of glucose levels: Jiangsu Nutrition Study

	Mortality Risk, HR (95% CI)		
	Deaths n (%)	Age and sex adjusted	Multivariable adjusted*
All-cause mortality			
NFG	146(5.7)	1.00	1.00
IFG	21(11.8)	1.54 (0.98 - 2.42)	1.83 (1.13 - 2.97)
Diabetes	17(16.7)	1.79 (1.09 - 2.96)	2.69 (1.62 - 4.49)
IFG/diabetes combined	38(13.6)	1.64(1.16 - 2.33)	2.12(1.44 - 3.12)
CVD mortality			
NFG	54(2.1)	1.00	1.00
IFG	9(5.1)	1.72 (0.82 - 3.63)	1.53 (0.70 - 3.37)
Diabetes	7(6.9)	2.09 (0.86 - 5.06)	1.97 (0.84 - 4.60)
IFG/diabetes combined	16(5.7)	1.87(1.02 - 3.41)	1.68(0.87 - 3.24)

*adjusted for age, gender, intake of energy, fat and fibre, smoking (0, 1-19, ≥ 20 cigarettes/day), alcohol drinking (no, 1-2 times/wk, 3-4 times/wk, daily), active commuting (no, 1-29 minutes/day, ≥ 30 minutes/day), leisure time physical activity (no, 1-29 minutes/day, > 30 minutes/day), sedentary activity (< 1 hrs/day, 1-1.9 hrs/day, 2-2.9 hrs/day, ≥ 3 hrs/day), education (low, medium, high), income (low, medium, high), occupation (manual/non-manual), BMI(linear and quadratic terms) and hypertension.

Figure 1 Age adjusted cumulative incidence of all-cause mortality according to glucose categories: the Jiangsu Nutrition Study.





Figure 2 Adjusted HR of IFG/diabetes (vs. NFG) for all-cause mortality according to sociodemographic and lifestyle factors.



Supplement Figure 1 Factor loadings of dietary patterns

